

PATIENTS' PRIORITIES AND ATTITUDES TOWARDS THEIR TEMPORO-MANDIBULAR DISORDERS

Martin Kelleher BDS MSc FDSRCPS FDSRCS, Consultant in Restorative Dentistry, King's College Dental Hospital, London

Arijit Ray-Chaudhuri BDS MFDS RCSEd MJDF RCSEng LLMA AHEA, Specialist Registrar in Restorative Dentistry, King's College Dental Hospital, London and St George's Hospital, London

Noman Khawaja BSc BDS MJDF RCSEng, Clinical Teacher in Primary Dental Care, King's College Dental Hospital, London

Keywords:

TMD, TMJ, temporo-mandibular joint, pain, survey, questionnaire, communication, history, management

Abstract

The diagnosis and appropriate management of temporo-mandibular disorders (TMDs) remains controversial. Current scientific evidence highlights the importance of psychosocial factors in sufferers and the reducing emphasis on occlusal or dental/prosthetic factors. This paper describes the findings of a survey of 211 patients reporting pain from their temporo-mandibular joint area and associated structures. This article offers busy primary dental care practitioners a cost effective questionnaire for obtaining relevant information from patients about the history of their condition and highlights what patients' hope to achieve through the management of their disorder. It also emphasises the importance of communicating effectively with patients and offers practical tips for the management of TMDs in primary care.

Introduction

In spite of decades of debate, discussion and dogma, the diagnosis and appropriate management of temporo-mandibular disorders (TMDs) remains highly contentious. Central to the problem that TMDs can pose for primary dental practitioners is that the term itself is poorly defined in the scientific literature and that its definitions have evolved over the last century.¹

The American Association of Dental Research (AADR) TMD Policy Statement Revision describes temporo-mandibular disorders as encompassing:

PATIENTS' PRIORITIES AND ATTITUDES TOWARDS THEIR TEMPORO-MANDIBULAR DISORDERS

"...a group of musculoskeletal and neuromuscular conditions that involve the temporo-mandibular joints, the masticatory muscles and all associated tissues. The signs and symptoms associated with these disorders are diverse and may include difficulties with chewing, speaking and other oro-facial functions".²

The World Health Organisation (WHO) also recognises that TMDs are separate conditions in their International Classification of Diseases and Related Health Problems (ICD-10). This classification recognises temporo-mandibular pain dysfunction syndrome, TMJ derangement, TMJ ligament strain and TMJ dislocation as different conditions affecting the same anatomical area.

Due to its complicated taxonomy, the epidemiological data are unclear about the prevalence and incidence of TMDs. The scientific evidence suggests that somewhere between 10% and 30% of the healthy adult population will be suffering current or recent symptoms of TMDs.^{3,4} This is a significant range but fortunately the vast majority of reported symptoms appear to be too mild and/or infrequent to trigger a request for professional help. Indeed it has been estimated that 85% of sufferers with signs and/or symptoms of TMD perceive that they have no treatment needs.⁵ The AADR TMD policy states that the differential diagnosis of these conditions and related oro-facial pains should be based on a thorough patient history and clinical examination in the first instance. It also recommends that other sophisticated TMJ investigations lack the requisite sensitivity and specificity to separate TMD sufferers from healthy patients. The vast majority of these electronic diagnostic devices, for example electromyography, have yet to be validated fully or scientifically and at present cannot be justified in the diagnosis of TMDs.⁶

Similarly, although several imaging modalities have been validated for their use in diagnosing TMDs their use is predominantly limited to the detection of intra-capsular pathology.⁷ The authors generally support the growing consensus that routine radiographic examinations such as dental panoramic radiographs or CBCT are of limited value in patients who present with TMDs, especially if this is of myogenic origin. These patients often do not have intra-capsular TMJ pathology and even when such pathology is present it is unlikely to be detected using such radiographic imaging. The commonest joint pathology found in TMD sufferers is related to the articular disc which is not visible with conventional radiography. If discal pathology is likely and requires further investigation then the authors recommend referral for Magnetic Resonance Imaging (MRI) and appropriate specialist reporting.

PATIENTS' PRIORITIES AND ATTITUDES TOWARDS THEIR TEMPORO-MANDIBULAR DISORDERS

History taking for patients with TMDs

There are several recognised and scientifically validated systems for recording the history of TMDs. The most commonly used is the Research Diagnostic Criteria for TMDs (RDC-TMD) which allows for a thorough exploration of both the clinical and behavioural aspects of the conditions.⁸ The RDC-TMD is used to provide epidemiological data for scientific research, but due to its complexity it is rarely appropriate for clinical use in primary dental care. For patients presenting with pain, or a history of pain, the authors recommend that a simplified pain history is obtained. Table 1 illustrates a commonly used structured questionnaire (SOCRATES) for obtaining a generic pain history.

An alternative history taking tool designed specifically for suspected TMD patients was used in this patient survey and is available from the corresponding author. Use of this written questionnaire avoids time consuming history taking, allowing the clinician to be focussed on the psychosocial aspects of the disorders as much as the clinical aspects. It also allows early identification of how the patient wishes to be managed and potentially avoids unnecessary or inappropriate interventions. What is important in obtaining any form of patient history is not what precise structure is used but rather that there is **some form of *structure* to the history taking**. If the patient is not suffering pain, they may alternatively describe one of the following common symptoms:

- A "click" when opening and/or closing their jaws
- A reduction of their ability to open or close their jaws comfortably
- Deviation on opening and/or closing of the jaws

PATIENTS' PRIORITIES AND ATTITUDES TOWARDS THEIR TEMPORO-MANDIBULAR DISORDERS

Heading	Question	Common answer(s)
Site	Where is the pain?	TMJ, muscles of mastication, one or both sides
Onset	When and how did the pain start?	More than 3 months ago, gradually
Character	What is the pain like e.g. sharp, dull, stabbing?	Dull, tenderness
Radiation	Does the pain spread anywhere?	Temple, ear, eye, one or both sides
Associations	Does anything else happen at the same time	Clicking of jaw joint, stiffness of jaw joint(s)
Timing	Does the pain follow any pattern?	Worse first thing in the morning, intermittent
Exacerbating/ Relieving factors	Does anything make the pain better or worse?	Better: painkillers, relaxation, holidays Worse: stress, eating, yawning
Severity	How bad is the pain out of 10?	Variable

Table 1: A pain history for TMDs using the 'SOCRATES' acronym. The right hand column provides common answers to these questions

Bruxism and tooth clenching

For many patients, their TMD is associated with periods of parafunctional oral habits. The commonest of these is teeth clenching or grinding (bruxism), which can happen during sleep as well as when the patient is awake. The scientific evidence for a direct relationship between clenching/bruxism and TMDs, however, is still weak. Several studies have identified a positive correlation, but some of these studies are characterised by methodological difficulties in relation to the correct identification of the two phenomena and bias.⁹

Interestingly, the WHO recognises teeth grinding (bruxism) in its ICD-10 and describes it as a 'somatoform disorder' under the "mental and behavioural disorders" sub-classification. It is classified by the WHO as being closely related to the following psychogenic conditions:

- Dysmenorrhoea – menstrual pain
- Dysphagia – difficulty swallowing
- Pruritus - itching
- Neck stiffness

PATIENTS' PRIORITIES AND ATTITUDES TOWARDS THEIR TEMPORO-MANDIBULAR DISORDERS

Grinding of teeth and the above problems are described together as:

"...disorders of sensation, function and behaviour, not due to physical disorders, which are not mediated through the autonomic nervous system, which are limited to specific systems or parts of the body, and which are closely associated in time with stressful events or problems"

Thus it is clear that the WHO define bruxism as a *psychogenic* condition and by inference not particularly amenable to correction through dental treatment.

However, many patients are not aware of or will not admit to any parafunctional habits that may be causing or contributing to their TMD, and will provide a negative history on questioning. They may, however, be aware of tiredness in their oro-facial musculature or some may describe an inability to locate their 'correct bite'. Clinical examination may also reveal signs which are not consistent with the absence of symptoms:

- Extraoral, eg. muscular hypertrophy, especially in the temporalis and masseter muscles
- Intraoral
 - Soft tissue changes eg. a white line in the cheek adjacent to the occlusal plane (linea alba), tongue scalloping or traumatic ulceration of the cheek or tongue
 - Hard tissue changes, eg. attritional tooth surface loss, cracked teeth or cracked/worn restorations

It is noteworthy that these findings may be in addition to, or exclusive of clinical findings of pain and/or dysfunction.

It is also worth noting that, as many patients are completely unaware of their TMDs and/or parafunctional habits, the dentist is often the first individual to piece together the various clues and provide a diagnosis. These clues should be carefully noted as the sub-clinical features of TMDs as parafunctional habits often impact on the dental management of the patient. This may be as simple as providing advice for the patient, fabrication of a removable appliance of varying designs, or the selection of an alternative material for a restoration. Alternatively, these diagnoses may have more far reaching implications such as whether to embark on or avoid complex restorative dental rehabilitation. Failure to obtain and document information relating to a patient's TMD and oral habits may leave the clinician vulnerable to a possible later medicolegal complaint.¹⁰

PATIENTS' PRIORITIES AND ATTITUDES TOWARDS THEIR TEMPORO-MANDIBULAR DISORDERS

Trends in the management of TMDs

In the last two decades, there has been a paradigm shift in the management of TMDs.¹¹ Modern management of TMDs has become far more focused on the identification of psychosocial factors and the patient's opinions and attitudes towards their management.¹² It was previously believed by some that bruxism and TMDs could be cured by dental or surgical interventions alone, without the need to explore the complex relationship between these two separate entities and the sufferer's psychosocial or social circumstances.^{13,14} This resulted in surgical procedures for the TMJ without recognition that myofacial and joint-related symptoms were often separate clinical entities. Similarly, earlier philosophies on dental management placed great emphasis on occlusal equilibration or extensive oral rehabilitation to provide the 'ideal' position of the mandibular condyles in the glenoid fossae, and this approach required significant irreversible treatment of the dentition. The authors' views are that if such destructive dental treatment is provided for a TMD patient who is later diagnosed to have a significant psychological component, the treating dentist could face a potential claim for negligence.¹⁵

A survey of patients reporting TMDs

Sadly, the scientific literature has been slow to reflect some of the modern management of TMDs. In addition, there is a dearth of reliable information in the UK on patients' subjective concerns, wishes or views on their temporo-mandibular pain or problems. The following section of this article seeks to help primary dental care clinicians by providing them with information on the demography of TMD patients as well as their likely priorities for the management of their condition. The authors recommend the use of the history taking questionnaire (available on request) as a practical, time saving tool for obtaining quantifiable evidence relating to their patients' reported symptoms and their specific treatment aims. This allows the management of TMDs to be based largely on the patient's own perceptions and priorities. The authors feel that this is a sensible pre-requisite for the successful modern management of TMDs in primary dental care.

Table 2 describes the demographic data of 211 consecutive patients with some symptoms or signs of TMD, mainly collected from general dental practitioners, who attended for a new patient consultation in the Department of Restorative Dentistry at Kings' College Hospital Dental Hospital,

PATIENTS' PRIORITIES AND ATTITUDES TOWARDS THEIR TEMPORO-MANDIBULAR DISORDERS

London between January 2008 and June 2009. This survey only included patients who agreed to complete a facial pain (FP) and hospital anxiety and depression (HAD) questionnaire and who were subsequently diagnosed with a TMD.

	Number	Percentage
Gender		
Female	165	78.2
Male	46	21.8
Marital status		
Unmarried	100	47.3
Married	72	34.1
Undisclosed	39	18.5
Education level		
Secondary up to 16	47	22.3
Secondary 18 or over	17	08.1
College/University	82	38.9
Undisclosed	65	30.8
Work status		
Unemployed	40	19.0
Employed	171	81.0

Table 2: A summary of the demographic data of the 211 patients who attended the Department of Restorative dentistry at King's College Hospital, London and were subsequently diagnosed with a TMD

Demographics

The patients ranged in age from 15 to 82 years with a mean of 39.2 years (SD 14.4). This broad range of ages in TMD sufferers is reported in several scientific studies, although subjective symptoms have been noted to reduce with age.^{3-5,16} In agreement with previous studies of TMDs, a significant majority of the surveyed patients were female.^{3-5,17,18} It was also interesting to note that the largest group of attending patients had entered higher education (38.9%). This was at variance with the patients who commonly attend the department, which is located in a deprived part of South London, UK. The majority of the patients diagnosed with a TMD were employed. The over-representation of patients with a higher socio-economic status attending for TMD diagnosis and management has been recognised for many years.¹⁶

PATIENTS' PRIORITIES AND ATTITUDES TOWARDS THEIR TEMPORO-MANDIBULAR DISORDERS

Affected activities reported by patients

Table 3 identifies the activities that were most likely to be affected by pain from the TMD. The figures illustrate that the activities that were affected were those that were most likely to directly require function of the TMJ and its associated structures, ie. chewing, eating hard foods and yawning. Activities that required less TMJ movement or muscular activity were far less likely to cause pain.

	Frequency	Number	Percentage
Chewing	1	168	89.4
Eating hard foods	2	154	82.8
Yawning	3	149	81.9
Smiling/Laughing	4	62	39.2
Talking	5	59	35.8
Eating soft foods	6	56	34.8
Drinking	7	35	22.0
Exercising	8	25	16.9
Swallowing	9	24	16.3
Other	10	21	41.2

Table 3: A summary of the activities that were most frequently affected by pain from a TMD

Patient expressed desires for management

Table 4 illustrates one of the most interesting results of this survey. The most frequently desired outcome (65%) by these patients was just to 'understand their pain better'. In addition, about 50% of patients wanted to 'know that their pain was not serious' while 21% of patients wished to 'improve communication with their clinician'.

PATIENTS' PRIORITIES AND ATTITUDES TOWARDS THEIR TEMPORO-MANDIBULAR DISORDERS

	Number	Percentage	Ranking
Understanding my pain problem more	137	65.0	1
Knowing pain is not serious	105	49.7	2
Able to eat with confidence	105	49.7	3
Returning or remaining at work	95	45.0	4
Reducing pain medication	62	29.4	5
Feel less self-conscious in public	57	27.0	6
Feeling less depressed	48	22.8	7
Improving communication with doctors about pain	65	20.8	8
Improving my ability to socialise	36	17.1	9
Being physically intimate with partner	31	14.7	10
Reduce tendency to overdo activities	26	12.3	11
Meeting others with similar pain	10	4.7	12

Table 4: A summary of the outcomes that patients identified as being 'very important' to them in their treatment for TMD

Discussion

Based on the AADR recommendations, scientific evidence and the results of this survey, the authors feel that the modern management of TMDs should begin with a detailed structured history of the patients' symptoms and parafunctional oral habits prior to examination for signs. An accurate medical history is an essential part of this, as sufferers are more likely to present with co-morbidities with overlapping symptoms with TMDs such as fibromyalgia, chronic fatigue syndrome, tension headaches, etc.¹⁷ The questionnaire included in this article allows for easy capture of this essential information and an early exploration of the patient's psychosocial background, as TMD patients are more likely to have suffered adverse life events, mood disorders, stress, anxiety and depression.^{11,18,19} This information, including their social history, should include a detailed analysis of any precipitating factors and recent life events such as changes in the workplace, ill health in the family, financial worries, bereavement etc. It is the opinion of the authors that it is not only negative life events that predispose patients to suffer TMDs, but also apparently positive ones such as positive changes in the patient's career or changes in their accommodation. It has long been recognised that all of these life events are predictors for future stress-related ill health.²⁰

PATIENTS' PRIORITIES AND ATTITUDES TOWARDS THEIR TEMPORO-MANDIBULAR DISORDERS

It is also beneficial to gauge the patient's desired outcomes at the time of the initial consultation. This allows individualisation of their management strategies and aids clinical decision-making with regard to the use of further imaging, fabrication of occlusal appliances of varying designs, or other specialist referral. The majority of the patients in this survey focused their desires on gaining more knowledge about their diagnosis and prognosis, rather than receiving active treatment. This correlates well with increasing calls for the management of TMDs to move from a surgical/dental-based model to a physician-based model.²¹ This approach allows clinicians to target their management on conservative strategies such as discussion of their disorder, explanation and reassurance, analgesic advice and/or occlusal appliance therapy (Table 5).

Read the questionnaire answers in advance of seeing the patient as background information
Listen empathetically
Provide a diagnosis of a TMD
Discuss pre-disposing factors to TMDs e.g. stress, depression, bruxism
Reassure the patient with regard to the benign and cyclical nature of the syndrome
Provide a written document in simple and appropriate language
Advise simple topical or oral NSAIDs
Advise application of warmth to the joint and/or musculature
Provide a full coverage splint if appropriate
Review the patient
Refer for further management if symptoms worsen or do not abate

Table 5: Key points in the conservative management of TMDs in primary dental care

Although the highest level of evidence for the efficacy of these conservative managements is not available, the authors of several scientific reviews conclude that due to their conservative and non-destructive nature, they should still be the mainstay of TMD management.²⁵⁻²⁸ This paucity of evidence at the systematic review level is more likely to be a reflection of a lack of scientific rigour rather than evidence of ineffectiveness. There is, however, a significant lack of high quality evidence relating to the invasive or speculative treatment of TMD with orthodontics, occlusal

PATIENTS' PRIORITIES AND ATTITUDES TOWARDS THEIR TEMPORO-MANDIBULAR DISORDERS

adjustment, arthrocentesis or injections of steroids, hyaluronate, other chemicals or open surgery of the TM joint itself. ²⁹⁻³²

The authors recognise that it may be difficult for busy primary care clinicians to devote a large amount of time to obtaining and recording this delicate information by the chair side. It is because of this that we recommend that a questionnaire similar to the one that was used in this study is sent to the patient ahead of the appointment. It can be completed in writing by the patient/partner/translator at home and in their own time prior to the consultation and then discussed at the chair side. This allows written exploration of a range of issues relevant to TMD which some dentists may find difficult to record in history taking eg. questions about anxiety and depression. It also allows patients to feel that their problems are being taken seriously and that they are not being forced to disclose delicate information at the chair side without prior warning. Finally, it can be stored in the patient's records as an excellent signed and dated summary of the patient's signs, symptoms and desires for the management of their condition.

Conclusions

The AADR and WHO use the term temporo-mandibular disorder to describe a heterogeneous group of pathologies affecting the TMJ and/or its associated musculature. The aetio-pathogenesis and clinical manifestations of TMDs are complicated and multi-faceted. This is often, although not always, associated with parafunctional oral habits such as clenching or bruxism. Patients diagnosed with the condition are also more likely to have recognised co-morbidities, precipitating psychosocial factors and/or mood affective disorders. This paper reports on the demographic data, activities affected and treatment desires of 211 TMD patients surveyed in an inner city hospital dental department.

The commonest demographic characteristics were that patients were likely to be

- Female
- Employed
- Well educated

The patients' activities that were mainly affected were:

PATIENTS' PRIORITIES AND ATTITUDES TOWARDS THEIR TEMPORO-MANDIBULAR DISORDERS

- Chewing
- Eating hard foods
- Yawning

The patients' main reasons for seeking care were:

- To gain more understanding of their pain problem
- To know that the pain was not serious
- To be able to eat again with confidence

A questionnaire has been developed which provides a simple and cost-effective way of helping to record, diagnose and discuss TMDs. Hopefully this will help clinicians to adopt a more modern approach to the management of TMDs which involving effective communication as well as being conservative and adopting reversible strategies in the first instance. To access the questionnaire, please contact arjit.ray-chaudhuri@nhs.net

Contact

For further information or to access the questionnaire, please contact Arijit Ray-Chaudhuri at aj10@hotmail.com

Acknowledgements

With acknowledgements to Dr. N.Donaldson at King's College Hospital, London, who gave valuable advice on the statistics.

References

1. Laskin DM. Temporomandibular disorders: a term past its time? JADA 2008;**139**:124-128
2. American Association of Dental Research. Policy statement: temporomandibular disorders 2010. American Association of Dental Research site. Available at: www.aadronline.org/i4a/pages/index.cfm?pageid=3465. Accessed June 12, 2012
3. Dworkin SF, Huggins KH, LeResche L, Von Korff M, Howard J, Truelove E et al. Epidemiology of signs and symptoms in temporomandibular disorders: clinical signs in cases and controls. JADA 1990;**120**:273-281
4. Goulet JP, Lavigne GJ, Lund JP. Jaw pain prevalence among French-speaking Canadians in Québec and related symptoms of temporomandibular disorders. J Dent Res 1995;**74**:1738-1744
5. De Kanter RJ, Truin GJ, Burgersdijk RC, Van 't Hop MA, Battistuzzi PGFCM, Kalsbeek H et al. Prevalence in the Dutch adult population and a meta-analysis of signs and symptoms of Temporomandibular disorder. J Dent Res 1993;**72**:1509-1518
6. Klasser GD, Okeson JP. The clinical usefulness of surface electromyography in the diagnosis and treatment of temporomandibular disorders JADA 2006;**137**:763-771
7. Makdissi. TMJ imaging: more pictures, less talk. FDJ 2011;**2**:172-182
8. Dworkin SF, Leresche L. Research diagnostic criteria for temporomandibular disorders: review, criteria, examinations and specifications, critique. J Craniomandib Disord 1992;**6**:301-355

PATIENTS' PRIORITIES AND ATTITUDES TOWARDS THEIR TEMPORO-MANDIBULAR DISORDERS

9. Manfredini D, Lobbezoo F. Relationship between bruxism and temporomandibular disorders: a systematic review of literature from 1998 to 2008. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2010;**109**:e26-50
10. Gray R, Al-Ani Z. Risk management in clinical practice. Part 8. Temporomandibular disorders.
11. *Br Dent J* 2010;**209**:433-449.
12. Greene CS, Klasser GD, Epstein JB. Revision of the American Association of Dental Research's Science Information Statement about Temporomandibular Disorders. *J Can Dent Assoc* 2010;**76**:a115
13. Rollman GB, Gillespie JM. The role of psychosocial factors in temporomandibular disorders. *Curr Rev Pain* 2000;**4**:71-81
14. Dimitroulis G. The role of surgery in the management of the TMJ: a critical review of the literature. Part 1. *Int J Oral Maxillofac Surg*. 2005;**34**:107-113
15. Dimitroulis G. The role of surgery in the management of the TMJ: a critical review of the literature. Part 2. *Int J Oral Maxillofac Surg* 2005; **34**:231-237
16. Schmitter M, Rammelsburg P, Hassel A. The prevalence of signs and symptoms of temporomandibular disorders in very old subjects. *J Oral Rehabil* 2005;**32**:467-473
17. Carlsson GE, Egermark I, Magnusson T. Predictors of signs and symptoms of temporomandibular disorders: a 20-year follow-up study from childhood to adulthood. *Acta Odontol Scand* 2002;**60**:180-185
18. Magnusson T, Egermark I, Carlsson GE. A longitudinal epidemiologic study of signs and symptoms of temporomandibular disorders from 15 to 35 years of age. *J Orofac Pain* 2000;**14**:310-319
19. Heloe B, Heloe LA. Characteristics of a group of patients with TMJ disorders. *Community Dent Oral Epidemiol* 1975;**3**:72-79
20. Aaron L, Burke M, Buchwald D. Overlapping conditions among patients with Chronic Fatigue Syndrome, Fibromyalgia and Temporomandibular disorder. *Arch Intern Med* 2000;**160**:221-227
21. Manfredini D. Mood and anxiety psychopathology and temporomandibular disorder: a spectrum approach. *J Oral Rehabil* 2004;**31**:933-940
22. Auerbach A, Laskin D, Frantsve LM, Orr T. Depression, pain, exposure to stressful life events and long-term outcomes in Temporomandibular disorder patients. *J Oral Maxillofac Surg* 2001;**59**:628-633
23. Holmes TH, Rahe RH. The social readjustment rating scale. *J Psychosom Res* 1967;**11**:213-218
24. Greene DM, Laskin CS. Temporomandibular Disorders: Moving from a Dentally Based to a Medically Based Model. *J Dent Res* 2000;**79**:1736-1739
25. Al-Ani MZ, Davies SJ, Gray RJ, Sloan P, Glenny AM. Stabilisation splint therapy for temporomandibular pain dysfunction syndrome. *Cochrane Database Syst Rev* 2004;**1**:CD002778
26. Macedo CR, Silva AB, Machado MAC, Saconato H, Prado GF. Occlusal splints for treating sleep bruxism (tooth grinding). *Cochrane Database Syst Rev* 2007;**4**:CD005514.
27. Aggarwal VR, Lovell K, Peters S, Javidi H, Joughin A, Goldthorpe J. Psychosocial interventions for the management of chronic orofacial pain *Cochrane Database Syst Rev* 2011;**11**:CD008456
28. Mujakperuo HR, Watson M, Morrison R, Macfarlane TV. Pharmacological interventions for pain in patients with temporomandibular disorders. *Cochrane Database Syst Rev*. 2010;**10**:CD004715
29. Luther F, Layton S, McDonald F. Orthodontics for treating temporomandibular joint (TMJ) disorders. *Cochrane Database Syst Rev* 2010;**7**:CD006541
30. Koh H, Robinson P. Occlusal adjustment for treating and preventing temporomandibular joint disorders. *Cochrane Database Syst Rev* 2003;**1**:CD003812
31. Guo C, Shi Z, Revington P. Arthrocentesis and lavage for treating temporomandibular joint disorders. *Cochrane Database Syst Rev* 2009;**4**:CD004973
32. Shi Z, Guo C, Awad M. Hyaluronate for temporomandibular joint disorders. *Cochrane Database Syst Rev* 2003;**1**:CD002970



PATIENTS' PRIORITIES AND ATTITUDES TOWARDS
THEIR TEMPORO-MANDIBULAR DISORDERS
